

Topological invariants of three-manifolds from $U_q(osp(1|2n))$

Sacha C. Blumen*

*School of Mathematics and Statistics,
University of Sydney, NSW 2006, Australia*

February 1, 2008

Abstract

We create Reshetikhin-Turaev topological invariants of closed orientable three-manifolds from the quantum supergroup $U_q(osp(1|2n))$ at certain even roots of unity. To construct the invariants we develop tensor product theorems for finite dimensional modules of $U_q(osp(1|2n))$ at roots of unity.

1 Introduction

Topological invariants of closed orientable three-manifolds may be constructed from modular or quasimodular Hopf algebras [1, 2]. Reshetikhin and Turaev's construction using modular Hopf algebras relies upon several theorems relating framed links in S^3 to closed orientable three-manifolds. The Lickorish-Wallace theorem states that each framed link in S^3 determines a closed, orientable 3-manifold and that every such 3-manifold is obtainable by performing surgery upon a framed link in S^3 . Kirby-Craggs, and Fenn and Rourke showed that homeomorphism classes of closed orientable three-manifolds may be generated by performing surgery upon elements of equivalence classes of framed links in S^3 , where the equivalence relations are generated by the Kirby moves. By taking such combinations of isotopy invariants of links in S^3 as to render them unchanged under the Kirby moves one obtains a topological invariant of 3-manifolds.

RT took invariants of isotopy derived from the quantum group $U_q(sl_2)$ at even roots of unity. Their method was adapted for the quantum algebras related to the A_n, B_n, C_n, D_n Lie algebras at all roots of unity [2, 3], and the exceptional quantum algebras and quantum superalgebras $U_q(osp(1|2))$ and $U_q(gl(2|1))$ at odd roots of unity [4]. Here we create invariants from $U_q(osp(1|2n))$ at $q = \exp(2\pi i/N)$ with $N = 2(2k+1)$.

*e-mail: sachab@maths.usyd.edu.au

Resethikhin-Turaev invariants can be constructed from a class of Hopf (super)algebras more general than (quasi)modular Hopf (super)algebras. For a Hopf (super)algebra A , invariants may be constructed if A has the following properties (where we take the quantum superdimension and quantum supertrace if A is a quantum superalgebra)

- i) there exists a finite collection of mutually non-isomorphic left A -modules $\{V_\lambda\}$ where λ ranges over some index set I such that $\dim(V_\lambda) < \infty$ and $\dim_q(V_\lambda) \neq 0, \forall \lambda \in I$,
- ii) for any finite collection of A -modules $V_{\lambda_1}, V_{\lambda_2}, \dots, V_{\lambda_s}$ such that $\lambda_i \in I$ for all i ,

$$V_{\lambda_1} \otimes V_{\lambda_2} \otimes \dots \otimes V_{\lambda_s} = \mathcal{V} \oplus \mathcal{Z}$$

where $\mathcal{V} = \bigoplus_{\lambda \in I} (V_\lambda)^{\oplus m(\lambda)}$, $m(\lambda) \geq 0$ is the multiplicity of V_λ in the direct sum, and $\mathcal{Z}$ is a possibly empty A -module with zero quantum dimension,

- iii) for each V_λ there is a dual module $(V_\lambda)^\dagger \cong (V_{\lambda^*})$ such that $\lambda^* \in I$ and there exists a distinguished module V_0 such that $(V_0)^\dagger \cong V_0$,
- iv) the central element $\delta = v - \sum_{\lambda \in I} d_\lambda \chi_\lambda(v^{-1}) C_\lambda$ vanishes upon acting on any V_μ where $\mu \in \Lambda_N^+$. Here $\chi_\zeta(v) = q^{-(\zeta, \zeta + 2\rho)}$ and $C_\lambda = \text{tr}_\lambda[(\text{id} \otimes \pi)(1 \otimes q^{2h_\rho}) R^T R]$. R is the universal R -matrix and $R^T = P.R.P$ where P is the permutation operator. $\{d_\lambda\}$ is a collection of complex valued constants such that at least one d_λ is non-zero. This condition ensures that combinations of isotopy invariants of links are unchanged under some of the Kirby moves, and
- v) the sum $z = \sum_{\lambda \in I} d_\lambda q^{-(\lambda, \lambda + 2\rho)} \dim_q(V_\lambda)$ is non-zero.

2 $U_q(\mathfrak{osp}(1|2n))$ at roots of unity and its finite dimensional modules

The quantum superalgebra $U_q(\mathfrak{osp}(1|2n))$ at roots of unity is not quasi-triangular. However, the quantum superalgebra now has a class of central elements which do not exist at generic q . These central elements generate an ideal of $U_q(\mathfrak{osp}(1|2n))$, which is also a two-sided co-ideal.

The quotient of $U_q(\mathfrak{osp}(1|2n))$ by this ideal turns out to be a quasi-triangular Hopf superalgebra. Let us denote this algebra by $U_q^{(N)}(\mathfrak{osp}(1|2n))$, the details of which may be found in [6].

Associated with $\mathfrak{osp}(1|2n)$ there is a euclidean space H^* which has a basis of vectors $\{\epsilon_i | 1 \leq i \leq n\}$ and a bilinear form such that $(\epsilon_i, \epsilon_j) = \delta_{i,j}$. The even (resp. odd) positive roots are $\Phi_0^+ = \{\epsilon_i \pm \epsilon_j, 2\epsilon_k | 1 \leq i < j \leq n, 1 \leq k \leq n\}$ (resp. $\Phi_1^+ = \{\epsilon_i | 1 \leq i \leq n\}$). Define $2\rho = \sum_{\alpha \in \Phi_0^+} \alpha - \sum_{\beta \in \Phi_1^+} \beta$, $\mathbb{Z} = \{\dots, -1, 0, 1, \dots\}$, $\mathbb{Z}_+ = \{0, 1, 2, \dots\}$ and $\mathbb{Z}_n = \{0, 1, 2, \dots, n-1\}$. Let $X \subseteq H^* = \sum_{i=1}^n \mathbb{Z}_+ \epsilon_i$ and $X_N = X/NX$.

Define $\phi_0 = \{\delta \in \Phi_0^+ \mid \delta/2 \notin \Phi_0^+\}$ and $\phi_1 = \Phi_1^+$. For a given $U_q^{(N)}(\mathfrak{osp}(1|2n))$ where $n \geq 2$ and $N \geq 3$ we define $\overline{\Lambda}_N^+$ by

$$\overline{\Lambda}_N^+ = \left\{ \lambda \in X \mid 0 \leq \frac{2(\lambda + \rho, \alpha)}{(\alpha, \alpha)} \leq N', \forall \alpha \in \phi \right\},$$

where $\phi = \phi_0 \cup \phi_1$ if $N \equiv 2 \pmod{4}$ and $\phi = \phi_0$ otherwise. Here $N' = N$ if N is odd and $N' = N/2$ otherwise. For $U_q^{(N)}(\mathfrak{osp}(1|2))$ where $N \geq 3$ we define $\overline{\Lambda}_N^+$ by

$$\overline{\Lambda}_N^+ = \{ \lambda \in X \mid 0 \leq (\lambda + \rho, \alpha) \leq N'' \},$$

where α is the single odd root and $N'' = N/4$ if $N \equiv 2 \pmod{4}$ and $N'' = N'$ otherwise. For each $\overline{\Lambda}_N^+$ we define Λ_N^+ identically except that we replace $\leq$ by $<$. Λ_N^+ plays the role of the index set I of the collection of $U_q^{(N)}(\mathfrak{osp}(1|2n))$ modules.

Let V be the fundamental module of $U_q^{(N)}(\mathfrak{osp}(1|2n))$ with highest weight ϵ_1 . V is irreducible, $2n + 1$ dimensional and has the same structure as the fundamental module of $U_q(\mathfrak{osp}(1|2n))$ [7]. In [7] we prove the following lemmas and theorems.

Lemma 2.1 *Set $N \geq 4$ to be even. For each $\mu \in \overline{\Lambda}_N^+$ there exists a finite dimensional $U_q^{(N)}(\mathfrak{osp}(1|2n))$ module V_μ with highest weight μ such that $V_\mu \subseteq V^{\otimes t}$ for some $t \geq 1$. The quantum superdimension of V_μ is $\text{sdim}_q(V_\mu) = \lim_{q \rightarrow 1} (\text{sdim}_q(V_\mu^{\text{gen}}))$ where V_μ^{gen} is the finite dimensional irreducible $U_q(\mathfrak{osp}(1|2n))$ module with highest weight μ . The $\text{sdim}_q(V_\mu) \neq 0$ for all $\mu \in \Lambda_N^+$ and $\text{sdim}_q(V_\mu) = 0$ for all $\mu \in \overline{\Lambda}_N^+ \setminus \Lambda_N^+$.*

Let W be the Weyl group of $\mathfrak{osp}(1|2n)$ and τ be the maximal element of W . Then $-\tau(\lambda) \in \Lambda_N^+$, $\forall \lambda \in \Lambda_N^+$, which implies that for each V_λ there is a dual module $(V_\lambda)^\dagger$ with highest weight in Λ_N^+ .

In theorems 2.1 and 2.2 we assume that $n \geq 2$ or that $n = 1$ and $N \neq 4, 8$.

Theorem 2.1 *Set $N \geq 4$ to be even, V to be the fundamental module of $U_q^{(N)}(\mathfrak{osp}(1|2n))$ and $\epsilon_1 \in \Lambda_N^+$. Then for every $t \in \mathbb{Z}_+$,*

$$V^{\otimes t} = \mathcal{V} \oplus \mathcal{Z}$$

where $\mathcal{V} = \bigoplus_{\lambda \in \Lambda_N^+} (V_\lambda)^{\oplus m(\lambda)}$, $m(\lambda) \in \mathbb{Z}_+$ and $\mathcal{Z}$ is a possibly empty direct sum of indecomposable modules, each with zero quantum superdimension.

Theorem 2.2 *Set $N \geq 4$ to be even and let V_{λ_i} be finite dimensional $U_q^{(N)}(\mathfrak{osp}(1|2n))$ modules with $\lambda_i \in \Lambda_N^+$ for all i . Then*

$$V_{\lambda_1} \otimes V_{\lambda_2} \otimes \cdots \otimes V_{\lambda_s} = \mathcal{V}' \oplus \mathcal{Z}'$$

where $s \geq 1$ and $\mathcal{V}'$ and $\mathcal{Z}'$ have the same form as in theorem 2.1

Lemma 2.2 *Set $N \geq 3$ to be odd, $\epsilon_1 \in \Lambda_N^+$ and $t \in \mathbb{Z}_{N/2+1/2-n}$. Then $V^{\otimes t} = \overline{\mathcal{V}}$ where $\overline{\mathcal{V}}$ has the same form as $\mathcal{V}$ does in theorem 2.1*

3 Finding the set $\{d_\lambda\}$

We know that properties i), ii) and iii) hold for $U_q^{(N)}(osp(1|2n))$ at even N . We now consider condition iv): there exists at least one collection of $\{d_\lambda\}$ that solves $\chi_\mu(v) = \sum_{\lambda \in \Lambda_N^+} d_\lambda \chi_\lambda(v^{-1}) \chi_\mu(C_\lambda)$ for all $\mu \in \Lambda_N^+$. Now at generic q , the eigenvalue of C_λ in an irreducible representation with highest weight $\mu \in \Lambda_N^+$ is given by $sch_\lambda(q^{2(\mu+\rho)}) = S_{\lambda,\mu}/Q_\mu$ where $S_{\lambda,\mu} = (-1)^{[\lambda]} \sum_{\sigma \in W} \epsilon'(\sigma) q^{2(\lambda+\rho, \sigma(\mu+\rho))}$, $Q_\mu = \sum_{\sigma \in W} \epsilon'(\sigma) q^{2(\rho, \sigma(\mu+\rho))}$, and where $\epsilon'(\sigma) = -1$ if the number of components of σ that are reflections with respect to the elements of ϕ_0 is odd and $\epsilon'(\sigma) = 1$ otherwise. Our proofs for theorems 2.1 and 2.2 tell us that $sch_\lambda(q^{2(\mu+\rho)})$ is well behaved when q is taken to the N^{th} root of unity, and yields the desired $\chi_\mu(C_\lambda)$. To simplify finding the $\{d_\lambda\}$ we initially consider $Q_\mu q^{-(\mu+2\rho, \mu)} = \sum_{\lambda \in \Lambda_N^+} d'_\lambda q^{(\lambda+2\rho, \lambda)} S'_{\lambda,\mu}$ where $d'_\lambda = (-1)^{[\lambda]} d_\lambda$ and $S'_{\lambda,\mu} = (-1)^{[\lambda]} S_{\lambda,\mu}$. To solve for the $\{d'_\lambda\}$ we consider

$$Q_\mu q^{-(\mu+2\rho, \mu)} = \sum_{\lambda \in X_N} x_\lambda q^{(\lambda+2\rho, \lambda)} S'_{\lambda,\mu},$$

and set $x_\lambda = cq^{-(\lambda, 2\rho)}$. We then have

$$Q_\mu q^{-(\mu+2\rho, \mu)} = \sum_{\sigma \in W} \epsilon'(\sigma) \sum_{\lambda \in X_N} cq^{(\lambda, \lambda)} q^{2(\lambda+\rho, \sigma(\mu+\rho))}.$$

To ensure x_λ is independent of μ we undertake the mapping $\lambda \rightarrow \sigma(\lambda + \rho) - \sigma(\mu + \rho)$ in the summation, which may be done as $\sigma(\lambda + \rho) - \sigma(\mu + \rho) \in X_N$ and the summation remains over X_N . We then obtain

$$Q_\mu q^{-(\mu, \mu+2\rho)} = \sum_{\lambda \in X_N} cq^{(\lambda, \lambda+2\rho) - (\mu, \mu+2\rho)} \sum_{\sigma \in W} \epsilon'(\sigma) q^{2(\rho, \sigma(\mu+\rho))}$$

which results in $c^{-1} = \sum_{\lambda \in X_N} q^{(\lambda, \lambda+2\rho)}$ and

$$x_\lambda = q^{-(\lambda, 2\rho)} / \sum_{\lambda \in X_N} q^{(\lambda, \lambda+2\rho)}.$$

Now $q^{(\lambda', \lambda'+2\rho)} = (-1)^p q^{(\lambda, \lambda+2\rho)}$ where $\lambda' = \lambda + N/2\epsilon_i$ for any ϵ_i and $p = 1$ if $N = 4k$, $p = 0$ if $N = 2(2k+1)$. Then c is not well defined if $N = 4k$ and we set $N = 2(2k+1)$ for the remainder of this paper. Now $x_{\lambda'} q^{(\lambda', \lambda'+2\rho)} S'_{\lambda', \mu} = x_\lambda q^{(\lambda, \lambda+2\rho)} S'_{\lambda, \mu}$ where $\lambda' = \lambda + N/2\epsilon_i$ for any ϵ_i . It then follows that $\sum_{\lambda \in X_N} x_\lambda q^{(\lambda+2\rho, \lambda)} S'_{\lambda, \mu} = 2^n \sum_{\lambda \in X_{N/2}} x_\lambda q^{(\lambda+2\rho, \lambda)} S'_{\lambda, \mu}$.

Let $\overline{N} = N/2$ and $\overline{\Lambda}_N^+$ be the fundamental domain for $X_{\overline{N}}$ under the action of the affine Weyl group $W_{\overline{N}}^{(N)}$ of $U_q^{(N)}(so_{2n+1})$ [3]. The affine Weyl groups $W_{N/2}$ of $U_q^{(N)}(osp(1|2n))$ and $U_q^{(N)}(so_{2n+1})$ are identical and as $\overline{\Lambda}_N^+ = \overline{\Lambda}_N^+$ it follows that $\overline{\Lambda}_N^+$ is a fundamental domain for $X_{N/2}$ under the action of $W_{N/2}$.

Now $S'_{\sigma(\lambda+\rho)-\rho,\mu} = \epsilon'(\sigma)S'_{\lambda,\mu}$ for any $\sigma \in W$. If $\lambda \in \overline{\Lambda}_N^+ \setminus \Lambda_N^+$ there either exists some $\sigma \in W$ such that $\sigma(\lambda + \rho) - \rho = \lambda$ and $\epsilon'(\sigma) = -1$ or some $w \in W$ such that $\epsilon'(w) = 1$ and $\lambda = w(\lambda + \rho) - \rho + kN/2\epsilon_i$ for some ϵ_i and $k \in \mathbb{Z}$. As $S'_{\lambda',\mu} = -S'_{\lambda,\mu}$ where $\lambda' = \lambda + N/2\epsilon_i$, it follows that $S'_{\lambda,\mu} = 0$ for $\lambda \in \overline{\Lambda}_N^+ \setminus \Lambda_N^+$. Then

$$2^n \sum_{\nu \in X_{N/2}} x_\nu q^{(\nu, \nu+2\rho)} S'_{\nu,\mu} = 2^n \sum_{\lambda \in \Lambda_N^+} \sum_{\sigma \in W} \epsilon'(\sigma) x_{\sigma(\lambda+\rho)-\rho} q^{(\lambda, \lambda+2\rho)} S'_{\lambda,\mu}.$$

As

$$\sum_{\lambda \in \Lambda_N^+} d'_\lambda q^{(\lambda+2\rho, \lambda)} S'_{\lambda,\mu} = 2^n \sum_{\lambda \in \Lambda_N^+} \sum_{\sigma \in W} \epsilon'(\sigma) x_{\sigma(\lambda+\rho)-\rho} q^{(\lambda, \lambda+2\rho)} S'_{\lambda,\mu}$$

we obtain

$$d_\lambda = q^{(2\rho, \rho)} \gamma \cdot \text{sdim}_q(V_\lambda) / \sum_{\mu \in X_{N/2}} q^{(\mu, \mu+2\rho)}$$

where γ is the denominator in the expression of the quantum superdimension. Note that $d_{\lambda^*} = d_\lambda$ where $\lambda^* = -\tau(\lambda)$. The denominator of the d_λ does not vanish: $\sum_{\mu \in X_{N/2}} q^{(\mu, \mu+2\rho)} = \sum_{\mu \in X_N} q^{(\mu, \mu+2\rho)} / 2^n$ and $\sum_{\mu \in X_N} q^{(\mu, \mu+2\rho)} = \prod_{k=0}^{n-1} G(N, 2k+1)$ where $G(N, m) = \sum_{i=0}^{N-1} q^{i(i+m)} = (1+i)\sqrt{N}/x^{m^2}$ and x is a complex primitive $4N^{\text{th}}$ root of unity.

4 Constructing the invariant

The final matter we need to consider is condition v): that

$$z = \sum_{\lambda \in \Lambda_N^+} d_\lambda q^{-(\lambda+2\rho, \lambda)} \text{sdim}_q(V_\lambda) \neq 0.$$

It follows from [2] that $z \neq 0$. Given $\zeta = \sum_{\lambda \in \Lambda_N^+} d_\lambda \text{sdim}_q(\lambda)$, $\zeta \neq 0$ and $z = d_0 \zeta$.

Now it is a relatively simple matter to construct the invariants. Denote a framed link in S^3 by L and the 3-manifold it gives rise to by M_L . Let $\sum(L)$ stand for the Reshetikhin-Turaev functor applied to L (see [1, 3, 4]). Set A_L to be the linking matrix of L defined by: a_{ii} is the framing number of the i^{th} component of L and a_{ij} , $i \neq j$ is the linking number between the i^{th} and j^{th} components of L . Let $\sigma(A_L)$ be the number of nonpositive eigenvalues of A_L . Then

$$\mathcal{F}(M_L) = z^{-\sigma(A_L)} \sum(L)$$

is a topological invariant of M_L .

References

- [1] Reshetikhin N Y and Turaev V G 1991 Invent. Math. 103 547-597

- [2] Turaev V G and Wenzl H 1993 Int. J. Math. 4 no. 2 323-358
- [3] Zhang R B 1996 Comm. Math. Phys. 182 no. 3 619-636
- [4] Zhang R B 1994 Mod. Phys. Lett. A 9 no. 16 1453-1465, 1995 Rev. Math. Phys. 7 no. 5 809-831, 1997 Lett. Math. Phys. 41 no. 1 1-11
- [5] Khoroshkin S M and Tolstoy V N 1991 Comm. Math. Phys. 141 no. 3 599-617
- [6] Zhang R B 1992 J. Math. Phys. 33 no. 11 3918-3930
- [7] Blumen S C “Tensor products of finite dimensional $U_q(osp(1|2n))$ modules at roots of unity”, to appear.